

## APLS: Cardiac Scenario 7

*This is a Teaching Scenario. Some flexibility in how it progresses is possible according to individual learner needs.*

### History {initial candidate briefing}

A 12-year-old male is admitted to the emergency department with a fractured femur after playing football.

#### **He has no other injuries and is previously well.**

To provide analgesia an initial dose of 1 microg/kg of I.V fentanyl is given, and a femoral nerve block is administered, using ropivacaine.

Shortly after performing the block the treating doctor is called urgently away. The child then has a brief grand mal convulsion and becomes apnoeic and pulseless. The nurse pushes the emergency bell and commences CPR. You are in ED and attend.

Estimated weight 40 kg.

### Initial impression {provide information as candidate assesses child and applies monitoring}

Apnoeic and pulseless. Nurse performing CPR.

### Clinical Course {to be given to candidate as they progress}

The child is in pulseless ventricular tachycardia (VT) which then changes to VF after the 1<sup>st</sup> shock. Following a 2<sup>nd</sup> shock, adrenaline and subsequent 2 minutes CPR, ROSC occurs with sinus rhythm 120/min and a palpable pulse. The child starts breathing and moving.

## INSTRUCTORS INFORMATION

### Key Treatment Points



<b>Airway &amp; Breathing</b>	Establish airway patency	
	BVM ventilation with 100% O <sub>2</sub>	
	Consider LMA/intubation or arrange for intubation	
<b>Circulation</b>	Pulseless VT & VF protocol	
<b>General Therapy</b>	Uninterrupted BLS	
<b>Specific therapy</b>	Lipid rescue, Intralipid 20%, 1.5 mls/kg and infusion. Ideally mentioned but not required for ROSC in the simulation	

**Diagnosis:** Pulseless ventricular tachycardia, VF, Ropivacaine (LA) toxicity

## Learning objectives

At the end of this session participants should be able to:

- Apply the structured approach to management and diagnosis during cardiac arrest
- Perform BLS/ALS effectively and safely
- Recall and apply the VF/VT ALS algorithm in their own practice
- Recall and apply the 4 Hs/Ts in their own practice

## Potential Issues to be Discussed

- Primary learning objectives are emphasis on VF/VT protocols
- Role of Lipid Rescue. If time for discussion. See RCH algorithm below

## Extra equipment

- Algorithm available in Monash Paediatric Emergency Medication Book
- Intralipid 20% 500 mls (Laminated card)

## Management of LA toxicity. RCH Guidelines

[https://www.rch.org.au/clinicalguide/guideline\\_index/Local\\_anaesthetic\\_poisoning/](https://www.rch.org.au/clinicalguide/guideline_index/Local_anaesthetic_poisoning/)

Modified for clarity. Algorithm available in Monash Paediatric Emergency Medication Book

### BLS/ALS protocols if indicated

**Immediately cease administration of the local anaesthetic. Call for help**

### Resuscitation

- Standard procedures and supportive care
- If there is evidence of cardiotoxicity, immediate BVM ventilation is necessary to prevent hypoxaemia, hypercarbia and acidosis. Prepare to intubate.
- Treat ventricular dysrhythmias with sodium bicarbonate (1–2 mmol/kg up to 100 mmol) IV; can use amiodarone; AVOID calcium channel blockers, beta blockers, local anaesthetics
- Treat seizures with benzodiazepines
- Treat hypotension with intravenous normal saline 10-20 mL/kg followed by inotropes if required; AVOID vasopressin
- Intravenous lipid emulsion (see below – antidotes)

### Antidotes

1. Sodium bicarbonate for ventricular dysrhythmias secondary to sodium channel blockade (see above).
2. Intravenous lipid emulsion (intralipid 20%) in severe cardiovascular toxicity or cardiac arrest.

Precautions - hypersensitivity to egg yolk, soya or peanut protein

### Administration of lipid emulsion therapy with 20% lipid emulsion

- Bolus 1.5 mL/kg IV over 1 minute
- Continuous infusion 0.25 mL/kg/minute. Wait 5 Minutes, then
  - Give a maximum of 2 repeat boluses for persistent cardiovascular collapse or deterioration (at least 5 minutes between boluses), AND
  - Double infusion to 0.5 mL/kg/minute
  - Continue infusion until stable and adequate circulation or maximum dose of lipid emulsion has been given (maximum cumulative dose is 12 mL/kg)

## APLS: Cardiac Scenario 8 Pacific

*This is a Teaching Scenario. Some flexibility in how it progresses is possible according to individual learner needs.*

### **History** {initial candidate briefing prior to arrival of child}

A 5 year old boy is brought to the Emergency department in the arms of his distressed parents. They say he was found unconscious in bed this morning having gone to sleep late the previous night with tummy ache.

Estimated weight 20 kg

### **Initial impression** {provide information as candidate assesses child and applies monitoring}

No response, apnoeic, pulseless.

### **Clinical Course** {to be given to candidate as they progress}

Initially in asystole: after ventilation with oxygen, chest compressions and one dose of adrenaline, the rhythm on the monitor remains asystole. After the 2<sup>nd</sup> cycle the rhythm is noted to be organised but there is still no pulse nor signs of life. Ventilation with oxygen and chest compressions continue and with a fluid bolus and a further dose of adrenaline the heart rate increases and a pulse can be felt.

### **Additional History and Observations**

Bruising of different colours on abdomen. Distension of abdomen.

## INSTRUCTORS INFORMATION

### Key Treatment Points



<b>Airway &amp; Breathing</b>	Establish airway patency	
	BVM ventilation with 100% O <sub>2</sub>	
	Consider LMA/intubation or arrange for intubation	
<b>Circulation</b>	Asystole then PEA protocol	
	Fluid bolus 10 ml/kg	
	IV/IO access	
<b>General Therapy</b>	Uninterrupted BLS	

**Diagnosis:** Asystole/PEA, non-accidental injury, sepsis – bowel perforation

### **Learning objectives**

At the end of this session participants should be able to:

- Apply the structured approach to management and diagnosis during cardiac arrest
- Perform BLS/ALS effectively and safely
- Recall and apply the ALS PEA and asystole algorithm in their own practice
- Recall and apply the 4 Hs/Ts in their own practice